

Math 320 Linear Algebra Assignment # 5

1. Let V be a vector space and $W = \{\vec{0}\}$. Show that W is a subspace of V . This is called the trivial subspace of V .
2. Let V be a vector space. And $\vec{v} \in V$ with $\vec{v} \neq \vec{0}$.
 - (a) Show if $r\vec{v} = \vec{0}$ then $r = 0$. (We showed the other direction in class).
 - (b) Show if $r_1, r_2 \in \mathbb{R}$ and $r_1\vec{v} = r_2\vec{v}$ then $r_1 = r_2$.
 - (c) Show that if W is a subspace of V and W is non-trivial (i.e. $W \neq \{\vec{0}\}$). then W is infinite.
3. For each of the following determine (with proof) if W is a subspace of the vector space V .

- (a) $V = \mathbb{R}^4$ and

$$W = \left\{ \begin{bmatrix} a + 2b \\ 0 \\ 3a + b \\ c \end{bmatrix} : a, b, c \in \mathbb{R} \right\}$$

- (b) $V = P_4$ and

$$W = \{ax^3 + bx^2 + 2x + c : a, b, c \in \mathbb{R}\}$$

- (c) $V = \mathbb{R}^{2 \times 2}$ and

$$W = \left\{ \begin{bmatrix} a & a^2 \\ b & 0 \end{bmatrix} : a, b \in \mathbb{R} \right\}$$

- (d) $V = \mathbb{R}^3$ and

$$\{\vec{x} \in \mathbb{R}^3 : A\vec{x} = \vec{b}\}$$

where:

$$A = \begin{bmatrix} 1 & -2 & 7 \\ 3 & -2 & -1 \\ -1 & 8 & 2 \end{bmatrix}, \quad \vec{b} = \begin{bmatrix} 2 \\ 1 \\ 4 \end{bmatrix}$$

- (e) $V = \mathcal{F}(\mathbb{R}, \mathbb{R})$ (the set of all function from the \mathbb{R} to the \mathbb{R}).

$$W = \{f \in \mathcal{F}(\mathbb{R}, \mathbb{R}) : f(2) = 0\}.$$

4. Let

$$A = \begin{bmatrix} 2 & 0 & 4 & 4 & 0 \\ 0 & 1 & -2 & -1 & 0 \\ 0 & 2 & -4 & -2 & 4 \\ 1 & 0 & 2 & 2 & 0 \end{bmatrix}$$

and

$$\vec{b} = \begin{bmatrix} 4 \\ 3 \\ 10 \\ 2 \end{bmatrix}$$

and

$$\vec{v}_1 = \begin{bmatrix} 2 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \quad \vec{v}_2 = \begin{bmatrix} 0 \\ 1 \\ 2 \\ 0 \end{bmatrix}, \quad \vec{v}_3 = \begin{bmatrix} 4 \\ -2 \\ -4 \\ 2 \end{bmatrix}, \quad \vec{v}_4 = \begin{bmatrix} 4 \\ -1 \\ -2 \\ 2 \end{bmatrix}, \quad \vec{v}_5 = \begin{bmatrix} 0 \\ 0 \\ 4 \\ 0 \end{bmatrix}$$

- (a) Does $A\vec{x} = \vec{b}$ have a solution? If so find one.
- (b) Is $\vec{b} \in \text{span}(\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4, \vec{v}_5)$? If so find $a_1, a_2, a_3, a_4, a_5 \in \mathbb{R}$ so that $b = a_1\vec{v}_1 + a_2\vec{v}_2 + a_3\vec{v}_3 + a_4\vec{v}_4 + a_5\vec{v}_5$. How is this related to the previous part?