

## Math 320 Linear Algebra Assignment # 10

1. Let

$$A = \begin{bmatrix} -2 & 1 & 0 & 0 & -1 \\ 3 & 2 & -2 & 0 & 0 \\ -3 & -1 & 3 & 5 & 1 \\ 0 & 0 & -2 & 0 & 3 \\ -3 & -1 & 1 & 0 & 0 \end{bmatrix}$$

- (a) Find  $\det(A)$  by expanding on the first row.
- (b) Find  $\det(A)$  by expanding on a different row or column.

2. Let  $A = \begin{bmatrix} 2 & 3 \\ -1 & 2 \end{bmatrix}$  and  $B = \begin{bmatrix} -3 & 1 \\ -2 & -2 \end{bmatrix}$ . Find the following:

- (a)  $\det(A)$
- (b)  $\det(B)$
- (c)  $AB$
- (d)  $\det(AB)$
- (e) Show  $\det(A)\det(B) = \det(AB)$ .

3. In this problem we are going to an important theorem. Let  $V$  and  $W$  be vector spaces and  $T : V \rightarrow W$  be an isomorphism (a 1-1 and onto linear transformation). Furthermore let  $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_r\} \subseteq V$  and  $\mathcal{D} = \{T(\vec{v}_1), T(\vec{v}_2), \dots, T(\vec{v}_r)\} \subseteq W$ .

- (a) If  $\mathcal{B}$  is linearly independent in  $V$  then  $\mathcal{D}$  is linearly independent in  $W$ . Here is a [short video](#) that will help with this part.
- (b) If  $\mathcal{B}$  spans  $V$  then  $\mathcal{D}$  spans  $W$ . Here is a [short video](#) that will help with this part.
- (c) If  $\mathcal{B}$  are a basis for  $V$  then  $\mathcal{D}$  are a basis for  $W$ .

(Hint: this should be very straight forward from the previous part)

4. Consider  $V = \mathbb{R}^{2 \times 2}$  and let

$$\mathcal{B} = \left\{ \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \right\}$$

- (a) What is  $\dim V$
- (b) Show that  $\mathcal{B}$  is linearly independent.
- (c) Why can you conclude from above that  $\mathcal{B}$  is a basis for  $V$ ?
- (d) Let

$$A = \begin{bmatrix} 2 & 4 \\ 1 & 1 \end{bmatrix}$$

Find  $[A]_{\mathcal{B}}$ . (Hint: Your answer should be an element of  $\mathbb{R}^4$ . Remember the order of  $\mathcal{B}$  matters.)

5. Let  $T : \mathbb{R}^{2 \times 2} \rightarrow P_2$  be defined by:

$$T \left( \begin{bmatrix} a & b \\ c & d \end{bmatrix} \right) = bx^2 + bx + c$$

(Hint: Only the first problem should require much in the way of work)

- (a) Find a basis for  $\ker(T)$ .
- (b) Find  $\dim(\ker(T))$ .
- (c) Find  $\dim(\operatorname{Rg}(T))$
- (d) Find a basis for  $\operatorname{Rg}(T)$
- (e) Is  $T$  onto?