

1. In class we defined Thomae's function $h : [0, 1] \rightarrow \mathbb{R}$ as:

$$h(x) = \begin{cases} 1 & \text{if } x = 0 \\ \frac{1}{n} & \text{if } x = \frac{m}{n} \text{ where } \gcd(m, n) = 1 \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

- (a) Show that h is not continuous for all $a \in \mathbb{Q}$.
- (b) Show that h is continuous for all $a \in \mathbb{R} \setminus \mathbb{Q}$.

2. Suppose that $[c, d] \subseteq [a, b]$. Remember that:

$$\mathbb{1}_{[c,d]}(x) = \begin{cases} 1 & \text{if } x \in [c, d] \\ 0 & \text{if } x \notin [c, d]. \end{cases}$$

Show that $\mathbb{1}_{[c,d]} \in \mathcal{R}[a, b]$ and that $\int_a^b \mathbb{1}_{[c,d]} = d - c$.

- 3. Suppose that $f \in \mathcal{C}([a, b])$, such that $f(x) \geq 0$ for all $x \in [a, b]$ and $\int_a^b f = 0$. Show $f(x) = 0$ for all $x \in [a, b]$
- 4. Find $f \in \mathcal{R}([a, b])$, such that $f(x) \geq 0$ for all $x \in [a, b]$ and $\int_a^b f = 0$, but $f(c) \neq 0$ for some $c \in \mathbb{R}$.
- 5. Let f be the fractional part function, i.e. $f(3.324) = 0.324$ ect.
 - (a) Graph f from 0 to 10
 - (b) Show that $f \in \mathcal{R}[0, 10]$.
 - (c) find $\int_0^{10} f$.