

The following problems are to get you ready for the final. You won't be turning them in so just use them to learn. Really try to solve them on your own and don't just watch someone solve them and think you can do it next time.

1. Give an example of each or explain why no such example exists.
 - (a) A non-continuous, non-monotone function on $[0, 1]$ that is in $R[0, 1]$.
 - (b) A function $f \in R[-1, 1]$ that has no anti-derivative.
 - (c) Bounded functions f and g defined on $[0, 1]$ with $f \in R[0, 1]$ and $g \notin R[0, 1]$ such that $f + g \notin R[0, 1]$.
 - (d) A function $f \in R[0, 1]$ such that $f(x) \geq 0$ for all $x \in [0, 1]$ and $\int_0^1 f = 0$ but $f(c) = 1$ for some $c \in [0, 1]$.
 - (e) A monotone function $f \in R[0, 1]$ such that $f(x) \geq 0$ for all $x \in [0, 1]$ and $\int_0^1 f = 0$ but $f(c) = 1$ for some $c \in (0, 1)$.
 - (f) A function $f \in C[0, 1]$ such that $f(x) \geq 0$ for all $x \in [0, 1]$ and $\int_0^1 f = 0$ but $f(c) = 1$ for some $c \in [0, 1]$.
 - (g) a function $f \in R[0, 1]$ that is neither monotone nor continuous
 - (h) a function $f \in R[0, 1]$ such that $|f| \notin R[0, 1]$
 - (i) a function $f \notin R[0, 1]$ such that $|f| \in R[0, 1]$
 - (j) a sequence of function f_n that converges to a function f pointwise on $[0, 1]$ but not uniformly on $[0, 1]$
 - (k) a sequence of function f_n all of which are not continuous, that converges uniformly on $[0, 1]$ to a function f that is continuous
 - (l) a power series with radius of convergence $R = 4$
 - (m) a power series that converges at $x = 2$ and $x = 4$ but not at $x = 0$
 - (n) a Hilbert space

2. What is the derivative of $F(x) = \int_x^{x^2} \ln(t) dt$ on $[1, e]$?

3. Suppose $f \in R[a, b]$ and $\int_a^b f = -1$. Also suppose there exists $d \in [a, b]$ such that $\int_a^d f = 3$. Show there exists $c \in [d, b]$ such that $\int_a^c f = 0$.

4. Define $f : [0, 1] \rightarrow \mathbb{R}$ by:

$$f(x) = \begin{cases} 1 & \text{if } x = \frac{1}{n} \text{ for } n \in \mathbb{N} \\ 0 & \text{otherwise} \end{cases}$$

Show $f \in R[0, 1]$ and compute the value of $\int_0^1 f$.

5. (a) Suppose $\{a_k\}$ and $\{b_k\}$ are sequences such that $|a_k| = |b_k|$ eventually, show that the power series $\sum_{k=0}^{\infty} a_k x^k$ and $\sum_{k=0}^{\infty} b_k x^k$ have the same radius of convergence.

(b) Use the previous part (and series you already know) to show that $\sum_{k=0}^{\infty} \frac{x^{2k}}{(2k)!}$ and $\sum_{k=0}^{\infty} \frac{x^{2k+1}}{(2k+1)!}$ both converge for all x .

(c) Define $\sinh(x) = \sum_{k=0}^{\infty} \frac{x^{2k+1}}{(2k+1)!}$ and $\cosh(x) = \sum_{k=0}^{\infty} \frac{x^{2k}}{(2k)!}$. Show $f_n(x) = \sum_{k=0}^n \frac{x^{2k+1}}{(2k+1)!}$ converges uniformly to $\sinh(x)$ on $[-N, N]$ for all $N \in \mathbb{N}$ and $g_n(x) = \sum_{k=0}^n \frac{x^{2k}}{(2k)!}$ converges uniformly to $\cosh(x)$ on $[-N, N]$ for all $N \in \mathbb{N}$.

(d) Prove $\frac{d}{dx} \sinh(x) = \cosh(x)$ and $\frac{d}{dx} \cosh(x) = \sinh(x)$.

(e) Let $H(x) = \sinh(x) + \cosh(x)$ show $H'(x) = H(x)$ and $H(0) = 1$. Thus it follows that $H(x) = \exp(x)$.

6. Suppose for all n , f_n is defined on $[0, 1]$ by:

$$f_n(x) = \begin{cases} ne^{-nx} & \text{if } x > 0 \\ 0 & \text{if } x = 0 \end{cases}$$

(a) Show $f_n \rightarrow f$ pointwise on $[0, 1]$ where $f(x) = 0$ for all $x \in [0, 1]$.

(b) Compute $\int_0^1 f_n$ for all n .

(c) Does $f_n \rightarrow f$ uniformly on $[0, 1]$?

7. Suppose V is a vector space and $\|\cdot\|$ is a norm on V . Let $a > 0$ and for all $v \in V$ let $\|v\|_a = a\|v\|$.

(a) Show $\|\cdot\|_a$ is a norm on V .

(b) Show that if d is the metric derived from $\|\cdot\|$ then show d_a defined by $d_a(x, y) = a[d(x, y)]$ for all $x, y \in V$ is the metric derived from $\|\cdot\|_a$.

(c) Show that $\{x_n\} \subseteq V$ and $x \in V$ then $x_n \rightarrow x$ using the metric d if and only if $x_n \rightarrow x$ using the metric d_a .

8. Find the radius for convergence of $\sum_{n=1}^{\infty} 3^n(x-2)^n$.

9. For all n let $f_n = \sin\left(\frac{x}{n}\right)$ defined on $[0, 1]$.

(a) Show $f_n \rightarrow 0$ pointwise on $[0, 1]$.

(b) Compute f'_n .

(c) Show that $f'_n \rightarrow 0$ uniformly on $[0, 1]$.

(d) Does $f_n \rightarrow 0$ uniformly on $[0, 1]$?

(e) Find $\int_0^1 f_n$.

(f) Use ?? to show that $\left[n - n \cos\left(\frac{1}{n}\right)\right] \rightarrow 0$.

10. Consider the metric space (\mathbb{R}, d) where:

$$d(x, y) = \begin{cases} 1 & \text{if } x = y \\ 0 & \text{otherwise} \end{cases} .$$

Does the sequence $\frac{1}{n}$ converge?

11. Let X and Y be normed vector spaces, and let $B(X, Y)$ be the space of bounded linear function from X to Y . In class, for $T \in B(X, Y)$ we defined:

$$\|T\|_{OP} = \sup \left\{ \frac{\|T(x)\|_Y}{\|x\|_X} : x \in X \right\} .$$

(a) Prove $\|\cdot\|_{OP}$ is a norm for $B(X, Y)$ (we did not ever show this in class).

(b) Prove that $B(X, Y)$ is a Banach space if Y is a Banach space.

12. Let $T : (L_2([1, 3]), \|\cdot\|_2) \rightarrow \mathbb{R}$ defined by $T(f) = 2 \int_1^3 f$.

(a) Show $T \in (L_2^*([1, 3]))$.

(b) Find (and prove) $\|T\|_{OP}$.

13. Let:

$$A = \begin{bmatrix} 2 & 3 \\ 0 & 2 \end{bmatrix} .$$

Find:

(a) $\|A\|_1$ (this is what we called $\|A\|_{11}$ before)

(b) $\|A\|_2$

(c) $\|A\|_u$