


IF
 RAMÓN Y CAJAL WERE ALIVE TODAY, WHICH REVIEW JOURNALS DO YOU THINK HE'D READ?

CHANCES ARE, YOU'D FIND THE LATEST ISSUE OF CURRENT OPINION IN NEUROBIOLOGY IN HIS LAP.







The inventor of tissue culture and first observer of live growth cones (1910) with the discoverer of the "organizer."



Hans Spemann & Ross Harrison

Figure 12.24 Growth Cone structure





Growth Cone extending *in vitro*

Mechanisms of Growth Cone Guidance

Contact-mediated (short-range)
- requiring direct cell-cell or cell-substrate contact

Contact attraction

Contact repulsion

Diffusible (' long-range')

Chemoattraction

Chemorepulsion

(Tessier-Lavigne & Goodman, 1996 "Molecular Biology of Axon Guidance")

Mechanisms of Growth Cone Guidance


Contact mediated (short-range) - requiring direct cell-cell or cell-substrate contact

Contact attraction - mediated by two classes of proteins

Direct Cell-Cell Adhesion: Cell Adhesion Molecules (CAMs)

Cell - Substrate Adhesion: Extracellular (ECM) Matrix proteins
and their cellular receptors

Figure 13.18 Outgrowth of Sensory Neurons
Cultured sensory neurons growing on stripes of laminin (ECM protein)



DEVELOPMENTAL BIOLOGY, Seventh Edition, Figure 13.18 © Garland Science, Inc. 2005. All rights reserved.

Mechanisms of Growth Cone Guidance

Cell Adhesion Molecules (CAMs)

Major classes:

Calcium-dependent Adhesion proteins: Cadherins
e.g. N-cadherin, E-cadherin, P-cadherin

Immunoglobulin Superfamily Adhesion proteins:
Ig-CAMs
e.g. NCAM, L1, Fasciclin II (FasII), etc.

Figure 3.28 Cadherin-Mediated Cell Adhesion




Figure 3.23 Reaggregation of Cells From Amphibian Neurulae





Fig 12.6 Expression of N- and E-cadherin Adhesion Proteins During Neurulation




Cadherins aid separation of epidermal and neural ectoderm

DEVELOPMENTAL BIOLOGY, Seventh Edition, Figure 12.6 (Part 1) © Garland Science, Inc. 2005 All rights reserved.


N-cadherin plays a role in neural crest cell migration

N-Cadherin level is high prior to neural crest cell migration, low during migration, and high again after migration ends.



Cell Adhesion

Cell adhesion can promote or inhibit migration



Mechanisms of Growth Cone Guidance

Cell Adhesion Molecules (CAMs)

Immunoglobulin Superfamily Adhesion proteins: Ig-CAMs
NCAM, L1, Fasciclin II, etc.

- NCAM - more general nervous system "glue"
 - used to direct retinal growth cone outgrowth (1 function)
 - two different forms based on glycosylation:
 - high SA (lower adhesion)
 - low SA (higher adhesion)
 - [SA = sialic acid]
 - high SA found during growth, low after reaching target

Mechanisms of Growth Cone Guidance

Cell Adhesion Molecules (CAMs)

Immunoglobulin Superfamily Adhesion proteins: Ig-CAMs
NCAM, L1, Fasciclin II, etc.

- L1 - more specific vertebrate CNS CAM
 - found in limited number of tracts, regions of brain & s.c.

Human mutations cause neurological disorders such as
X-linked hydrocephalus, MASA syndrome

L1 disruption by ethanol may cause part of Fetal Alcohol
Syndrome birth defects

Figure 21.13(2) Possible Mechanisms Producing Fetal Alcohol Syndrome (FAS)






Figure 21.12 Comparison of a Brain from an Infant with Fetal Alcohol Syndrome (FAS, Left) With a Brain From a Normal Infant of the Same Age (Right)



DEVELOPMENTAL BIOLOGY, Seventh Edition, Figure 21.12 © Garland Science, Inc. © 2005 All rights reserved.

Figure 21.13(1) Possible Mechanisms Producing Fetal Alcohol Syndrome (FAS)



Nile-Blue stained dying cells in 9d mouse embryo brain


DEVELOPMENTAL BIOLOGY, Seventh Edition, Figure 21.13 (Part 1) © Garland Science, Inc. © 2005 All rights reserved.


Mechanisms of Growth Cone Guidance


Contact attraction by CAMs

The "Labeled Pathways" Hypothesis -

Most growth cones are guided along axons already laid down (by "pioneer neurons") - each with a unique molecular marker or set of markers







A Grid of Pioneer Axon Bundles Guides Later Growth Cones, such as 'G'

Experiments:

Kill cells that make A/P bundle:
G growth cone guidance disrupted

Block function of Fas II (found on A/P bundle):
G growth cone guidance disrupted

Mechanisms of Growth Cone Guidance

Contact attraction by CAMs

Intermediate cell recognition by pioneer neurons:
"guidepost cells" in grasshopper limb bud

Pioneer Neurons in the Grasshopper Limb Bud







Fig. 13.20 The Action of Semaphorin 1 in the Developing Grasshopper Limb




Crossing of retinal axons at the optic chiasm



Ephrins act in contact repulsion at the optic chiasm






Mechanisms of Growth Cone Guidance

Extracellular Matrix proteins:

Collagen, Fibronectin, Laminin, etc.

ECM Receptors:

Integrins (alpha and beta subunits)



Mechanisms of Growth Cone Guidance

Chemoattraction by diffusible substances

Chemoattraction of growth cones first shown with NGF




Fig 13.27 Embryonic Axon From a Rat Dorsal Root Ganglion Turn in Response to NT-3




Figure 13.22 Trajectory of the Commissural Axons in the Rat Spinal Cord




Fig. 13.23 Transformed Chick Fibroblast (COS) Cells Secreting Netrins Elicit Axon Outgrowth of Commissural Neurons from 11-d Emb. Rat Dorsal Spinal Cord Explants





Fig. 13.23 Transformed Chick Fibroblast (COS) Cells Secreting Netrins Elicit Axon Outgrowth of Commissural Neurons from 11-d Emb. Rat Dorsal Spinal Cord Explants



Signals can be attractive or repulsive, depending on the receptor

Netrin-mediated dorsal/ventral signaling is an ancient feature of animal nervous systems.

Netrin signaling also patterns growth cone guidance and cell migration in the nematode *C. elegans*.

Mutants with abnormal nervous system function ("unc" - uncoordinated) were found to be affected in netrin-mediated growth cone guidance.

The first example of a netrin receptor mediating *chemorepulsion* was discovered in *C. elegans* (the *unc-5* gene).

Chemorepulsion by netrins is also found in the vertebrates.

Figure 13.24 C.e. Netrin Expression And Function in Axonal Guidance






Figure 13.25 Netrins inhibit Outgrowth of Trochlear Axons From Explants of the Mouse Brain Stem




A vertebrate example of Chemorepulsion by a netrin


DEVELOPMENTAL BIOLOGY, Seventh Edition, Figure 13.25 © Garland Science, Inc. © 2005. All rights reserved.





Neuronal Development


Neuronal cell death is a normal part of development in all nervous systems.


















Discoverers of Nerve Growth Factor (NGF)




Viktor Hamburger



Rita Levi-Montalcini

Stanley Cohen and Rita Levi-Montalcini awarded Nobel Prize for their work on NGF in 1986


NGF function is essential for neuronal survival *in vivo*



Levi-Montalcini, 1972

Neurotrophic Factors support the survival of specific types of neurons

Neurotrophins - family of NGF-related proteins
NGF - Nerve Growth Factor
BDNF - Brain-derived neurotrophic factor
NT3 - Neurotrophin 3
NT4 - Neurotrophin 4



Neurotrophic Factors support the survival of specific types of neurons

Neurotrophins - family of NGF-related proteins
NGF - Nerve Growth Factor
BDNF - Brain-derived neurotrophic factor
NT-3 - Neurotrophin 3
NT-4 - Neurotrophin 4

13 kDa secreted glycoproteins (as homodimer)

Bind to receptor tyrosine kinases: TrkA, TrkB, TrkC

e.g., NGF receptors - TrkA

BDNF, NT-4 receptor - TrkB

NT-3 receptor - TrkC

(NT-3 also binds to TrkA, TrkB)

Neurotrophic Factors support the survival of specific types of neurons


Neurotrophins - family of NGF-related proteins
NGF - Nerve Growth Factor
BDNF - Brain-derived neurotrophic factor
NT3 - Neurotrophin 3
NT4/5 - Neurotrophin 4/5

Other Neurotrophic and Differentiation Factors:

GDNF- Glial-derived neurotrophic factor

CDF/LIF - Cholinergic differentiation factor/
Leukemia inhibiting factor

Figure 13.29 Effects of NGF And BDNF On Axonal Outgrowth



DEVELOPMENTAL BIOLOGY, Seventh Edition, Figure 13.29 © 2005 Sinauer Associates, Inc. © 2005 All rights reserved.

Neurotrophic Factors support the survival of specific types of neurons

Three potential functions of neurotrophic and related factors


1. *Neurotrophic* factor
2. *Chemoattractive* factor
3. *Instructive* (inductive/signaling) factor

NGF shows all three functions

Experiment showing NGF's instructive role:

NGF can convert presumptive adrenal medulla cells (would become chromaffin cells) into neurons.

Figure 13.7 Differentiation of trunk neural crest cells



Neuronal Differentiation Factors

Example: CDF/LIF

CDF is made by heart muscle cells.

Some NC-derived neurons that innervate heart initially make norepinephrine (NE).

Interaction with heart converts cells to use ACh.

CDF has other roles in development:

LIF (immune system)

Essential for implantation of blastocyst
